

EXAM 1

EECS 203

Spring 2015

Name (Print): _____

uniquname (Print): _____

Instructions. You have 110 minutes to complete this exam. You may have one page of notes (8.5x11.5 two-sided) but may not use any other sources of information, including electronic devices, textbooks, or notes. Leave at least one seat between yourself and other students. Please write clearly. If we cannot read your writing, it will not be graded.

Honor Code. This course operates under the rules of the College of Engineering Honor Code. Your signature endorses the pledge below. After you finish your exam, please sign on the line below:

I have neither given nor received aid on this examination, nor have I concealed any violations of the Honor Code.

Page #	Points
2&3	/21
4	/10
5	/7
6	/10
7	/12
8	/10
9	/8
10	/10
11	/12
Total	/100

1. Logic and sets (21 points)

In this section, each question will have zero or more correct answers. You are to circle each correct answer and leave uncircled each incorrect answer.

[3 points each, -1 per incorrect circle/non-circle, minimum 0 points per problem]

a) Let w , b and n be propositions where

- w is "I walk to work"
- b is "I work at Burger King"
- n is "I work at night"

The sentence "When I work nights and I work at Burger King, I don't walk to work" could be written using propositions and logical connectives as:

$$(n \wedge b) \rightarrow \neg w$$

$$(n \vee b) \leftrightarrow n$$

$$n \rightarrow \neg (w \wedge b)$$

$$\neg(w \wedge b) \vee n$$


b) The circuit on the performs an operation that is equivalent to:

$$(\neg p \wedge q) \vee \neg p$$

$$\neg((\neg p \wedge q) \vee p)$$

$$\neg q$$

$$\neg(p \vee q)$$



c) Circle each of the following that is a *tautology*.

$$(a \rightarrow b) \leftrightarrow (\neg a \rightarrow \neg b)$$

$$(a \rightarrow b) \rightarrow (\neg b \rightarrow \neg a)$$

$$(a \wedge b) \rightarrow a$$

$$(a \wedge b \wedge c) \leftrightarrow (a \wedge c \wedge b)$$

d) Circle each of the following that is *satisfiable*.

$$(a \vee b) \wedge (a \vee \neg b) \wedge (\neg a \vee b) \wedge (\neg a \vee \neg b)$$

$$(a \rightarrow b) \rightarrow (\neg b \rightarrow \neg a)$$

$$(a \wedge b) \wedge (a \wedge \neg b)$$

$$(a \wedge b) \leftrightarrow (a \wedge \neg b)$$

e) We define the following predicates:

- $C(x)$: x is a car
- $F(x,y)$: x is faster than y
- $B(x)$: x has brakes

Consider the statement "Any car with brakes is faster than at least one car that doesn't have brakes."

Zero or more of the expressions below are accurate translations of this statement. Circle each of the following that are correct.

$$\exists x \forall y (C(x) \wedge F(y, x) \wedge B(y))$$

$$\forall y \exists x (C(x) \wedge F(x, y) \wedge B(y))$$

$$\forall x \exists y (C(x) \wedge C(y) \rightarrow B(y) \wedge F(x, y))$$

$$\forall x \exists y (C(x) \wedge B(x) \rightarrow C(y) \wedge \neg B(y) \wedge F(x, y))$$

$$\forall x (C(x) \wedge B(x)) \rightarrow \forall x \exists y (C(y) \wedge \neg B(y) \wedge F(x, y))$$

f) Circle each of the following which are *tautologies*.

$$(A \subset (A \cup B)) \rightarrow (B \subset A)$$

$$|A \cup B| \geq |A| + |B|$$

$$|A \cap B| \leq |A| + |B|$$

$$(B \subset A) \rightarrow (A \subset (A \cup B))$$

$$(A - B = \{1\}) \rightarrow ((1 \in A) \vee (1 \notin B))$$

g) Circle each of the following which are *tautologies* (note: ϕ is the empty set)

$$\phi \subset \{\{\phi\}\}$$

$$\{\phi\} \subset \{\{\phi\}, \phi\}$$

$$\phi \cap A \subseteq \phi$$


$$\phi \cup A \supset \phi$$

$$\{\phi\} \subset \{1, 2\}$$

2. More Sets (4 points)

(No partial credit will be given on this problem.)

Shade the Venn diagram below to show $(B - A) \cup (\overline{C} \cap \overline{B}) \cup (B \cap C)$



3. Functions (6 points)

In this section, each question will have zero or more correct answers. You are to circle each correct answer and leave uncircled each incorrect answer.

[3 points each, -1 per incorrect circle/non-circle, minimum 0 points per problem]

- a) Say that f is a function from $A \rightarrow B$ where A and B are both subsets of \mathbb{N} (the natural numbers). If $|A| > |B|$ then you can conclude that

f is not onto

f is not one-to-one

f is a bijection

f is not a bijection

The cardinality of B is greater than or equal to the cardinality of the range of f .

- b) Say that f is a function from $A \rightarrow B$ where A and B are both subsets of \mathbb{N} (the natural numbers). If $A \subset B$ then you can conclude that

f is not onto

f is not one-to-one

f is a bijection

f is not a bijection

The cardinality of B is greater than or equal to the cardinality of the range of f .

4. More functions (7 points)

Each part has one correct answer. **(-2 for each wrong or blank answer, minimum 0)**

For each of the following mappings indicate what type of function they are (if any). Use the following key:

- i. Not a function
- ii. A function which is neither onto nor one-to-one
- iii. A function which is onto but not one-to-one
- iv. A function which is one-to-one but not onto
- v. A function which is both onto and one-to-one

a) The mapping f from \mathbb{R} to \mathbb{R} defined by $f(x) = -2x$.

- i ii iii iv v

b) The mapping f from \mathbb{Z}^+ to \mathbb{Z}^+ defined by $f(n) = n - 1$.

- i ii iii iv v

c) The mapping f from \mathbb{Z} to \mathbb{Z} defined by $f(n) = 2n + 1$.

- i ii iii iv v

d) The mapping f from \mathbb{Z} to \mathbb{Z} defined by $f(n) = n + 1$.

- i ii iii iv v

e) The mapping f from \mathbb{R} to \mathbb{Z} defined by $f(x) = [x]$.

- i ii iii iv v

f) The mapping f from \mathbb{Z} to \mathbb{Z} defined by $f(x) = \frac{1}{x^2}$.

- i ii iii iv v

5. Growth of functions and infinite sets (10 points)

In this section, each question will have zero or more correct answers. You are to circle each correct answer and leave uncircled each incorrect answer.

Each problem is worth 2 points and you only get the points if you circle all of the correct answers and none of the wrong ones.

a) $12X^2\log(X)+X$ is:

$\Theta(X^2)$ $O(X^2)$ $\Omega(X^2)$ $\Theta(X^3)$ $O(X^3)$

b) Consider the following pseudo code:

```
for (i:=1 to 3)
  for (j:= 1 to n)
    if (A[i]>A[j])
      swap(A[i],A[j]); //Takes  $\Theta(1)$  time.
```

This algorithm has a run time of

$\Theta(n)$ $\Theta(n^2)$ $\Omega(\log(n))$ $O(n^3)$ $O(n/2+1)$

c) $X^4+12X^2\log(X)+X$ is:

$\Theta(X^4)$ $O(X^4)$ $\Omega(X^4)$ $\Theta(X^3)$ $\Omega(X^3)$

d) If A and B are both countably infinite sets, then $A \cap B$ could be

Countably infinite **Uncountably infinite** **Finite**

e) If A and B are both uncountably infinite sets, then $A \cup B$ could be

Countably infinite **Uncountably infinite** **Finite**

6. Number theory questions (12 points)

Provide your answers below and provide work when requested. Partial credit will not be given for incorrect answers (though it might be if you get the right answer without clear work where it is required).

- a. Compute $7^{11} \bmod 10$. Show your work. **[3]**

- b. Convert 1001011101_2 to base 16. **[2]**

- c. How many zeros are at the end of $50!$? (50 factorial) Show your work. **[3]**

- d. What is $\gcd(5454, 2700)$? **[2]**

- e. Which of the positive integers less than 9 are relatively prime to 9? **[2]**

7. Proof by induction (10 points)

Use induction to prove that $\sum_{k=1}^n k^2$ is equal to $\frac{n(n+1)(2n+1)}{6}$

Theorem:

Proof:

Base case:

Induction step:

8. Deductive proofs (8 points)

Provide a deductive proof to show that from the premises

- $(p \wedge q) \rightarrow r$
- $\neg r \rightarrow (p \wedge q)$
- $p \vee q$
- $r \rightarrow \neg q$

That we can conclude

- p

9. Rules of logic (10 points)

Prove that $(\neg p \rightarrow ((q \vee r) \wedge (\neg q \wedge \neg r))) \rightarrow p$ is a tautology using the rules of logic.

10. Other techniques (12 points)

a) Redo problem 8 but prove it using truth tables. **[6 points]**

b) Find a bijection from $[0,1]$ to $(0,1)$. **[6 points]**