EECS 206 – Winter 2002

Homework #6 – Assigned Feb. 15 – Due Friday Feb. 22

Relevant Lectures: 2/11, 2/13, 2/15

Relevant Reading: 4.1 (up to the end of 4.1.1), 9.1.1, 9.3.1, 9.3.2, Section 3 in G. Wakefield's "Primer on DFT", Section 4.2.3 in Lab. 4 handout.

Relevant Items in the DSP First CD: None this week.

Homework Submission Policies: Same as before (see course Web page).

- 1. Calculate the N-point DFTs of the following signals:
 - (a) s[n] = 1 for n = 0, s[n] = 0 for $1 \le n \le N 1$
 - (b) s[n] = 1 for $0 \le n \le N 1$
 - (c) $s[n] = \sin(2\pi mn/N)$ for $0 \le n \le N-1$. *Hint: See your lecture notes for 2/15!*
- 2. Suppose that the DFT of a periodic signal x[n] is X[0] = 1 and X[k] = 0 for all other values of k. What is x[n]?
- 3. Suppose that we have a real signal x[n], with fundamental period N_0 , whose DFT coefficients are all equal to zero except X[k] and $X[N_0 k]$. Let $X[k] = Ae^{j\phi}$.
 - (a) What is the average value of x[n]?
 - (b) Find x[n]. *Hint: Find* $X[N_0 - k]$ *first.*
 - (c) What is the average power of x[n]?
- 4. Consider the synthesis equation of the DFT:

$$x[n] = \sum_{k=0}^{N_0 - 1} X[k] e^{j\frac{2\pi}{N_0}kn} .$$

Use that equation to show that $x[n - N_0] = x[n]$.

5. Prove the time-shift DFT property:

$$y[n] = x[n - n_1] \Longrightarrow Y[k] = e^{-j\frac{2\pi}{N_0}kn_1}X[k]$$

- 6. Calculate the 4-point DFTs of the following 4-point signals:
 - (a) $x[n] = \cos(\pi n/2), n = 0, 1, 2, 3$
 - (b) $x[n] = 2^n, n = 0, 1, 2, 3$

7. Consider the discrete-time square wave of fundamental period N_0 , where it is assumed that N_0 is even:

$$x[n] = \begin{cases} 1 & 0 \le n \le \frac{N_0}{2} - 1\\ 0 & \frac{N_0}{2} \le n \le N_0 - 1 \end{cases}$$

- (a) Calculate X[0]
- (b) Verify that for $k \neq 0$,

$$X[k] = e^{-j\frac{2\pi}{N_0}k\frac{1}{2}(\frac{N_0}{2}-1)}\frac{\sin(\frac{2\pi}{N_0}k\frac{N_0}{4})}{\sin(\frac{\pi k}{N_0})}$$

Hint: After writing the DFT-analysis equation, use the fact below about geometric series and then factor the first term in the given expression for X[k]; the sin terms should "appear"...

$$\sum_{n=0}^{N-1} \alpha^n = \frac{1-\alpha^N}{1-\alpha} \quad \text{whenever } \alpha \neq 1$$