

A LINEAR ALGEBRAIC FRAMEWORK FOR THE ANALYSIS OF DISCRETE-TIME NONLINEAR SYSTEMS*

J. W. GRIZZLE†

Abstract. A linear algebraic framework for the analysis of synthesis-type problems for discrete-time nonlinear systems is introduced. This is an extension of a similar tool for continuous-time systems that established important connections between many algorithms associated with right-invertibility, left-invertibility and dynamic decoupling, as well as between these algorithms and an approach based upon differential algebra. A similar payoff is seen to be possible in the discrete-time setting.

Key words. nonlinear systems, discrete-time, ranks, invertibility

AMS subject classifications. 93C10, 49E05

1. Introduction. This paper extends to the class of discrete-time nonlinear systems the linear algebraic framework of [4], which has proven useful in the analysis of several synthesis problems for the class of continuous-time nonlinear systems [1]–[3], [13], [16], [32]. Recall that [4], through the introduction of a chain of subspaces naturally associated with the output of a system, provided a high-level interpretation of the inversion and dynamic decoupling algorithms that are built around the recursive computation of certain ranks associated with left-invertibility, right-invertibility, and noninteracting control. In addition, it established relationships between these algorithms and the differential algebraic approach. This same linear algebraic setting has been used in [3] to formulate in an intrinsic way the regularity (constant rank) conditions common to several procedures for synthesizing nonlinear dynamic compensators.

The reader is reminded that the importance of algebraic techniques and reasoning for analyzing many aspects of discrete-time nonlinear systems has been firmly established in [8]–[12], [25], [28], [29], and the references therein. The techniques employed here are most closely related to those of [10] and [28].

When studying continuous-time nonlinear systems, the class of affine systems (so-called because the dynamics is affinely parametrized by the control variables) has received the bulk of the attention of the nonlinear community. This is true for several reasons, the most important of which is that the class of affine systems is general enough to encompass many models arising in practice. However, it is also specific enough to admit reasonably simple analyses from at least two perspectives: *geometrically*, we are working with a finite number of vector fields, a drift term, and m control vector fields, as opposed to some arbitrarily, smoothly parametrized family of vector fields; *algebraically*, the various derivatives of the outputs depend polynomially on the inputs and their derivatives (with coefficients depending on the state, and the highest order derivative of the input appearing affinely), as opposed to some more general nonlinear dependence on the input. However, if we accept as an axiom that

* Received by the editors May 13, 1991; accepted for publication (in revised form) March 23, 1992. This work was completed while the author was a visiting professor (Poste Rouge, Centre National de la Recherche Scientifique) at the Laboratoire des Signaux et Systèmes, École Supérieure d'Électricité-Centre National de la Recherche Scientifique, Gif-sur-Yvette, France. This research was supported by the National Science Foundation contract NSF ECS-88-96136.

† Department of Electrical Engineering and Computer Science, University of Michigan, Ann Arbor, Michigan 48109-2122.

any interesting class of discrete-time systems should include time-sampled (digital) versions of the class of continuous-time affine systems, then we are obliged to consider systems of the form

$$\begin{aligned}x[k+1] &= f(x[k], u[k]), \\ y[k] &= h(x[k], u[k]),\end{aligned}$$

where f and h are sufficiently smooth functions, but otherwise arbitrary (consider sampling a continuous-time bilinear system). Consequently, it is not possible to assume that the dynamics is affine in the control variable (and hence, finitely parametrized); and even if it were, this would not entail that the iterates of the output depend polynomially on the inputs, with the highest-order delayed input appearing affinely. Consequently, the proof techniques of [4], based upon "global" interpretations of the inversion and dynamic extension algorithms, cannot be easily extended to discrete-time systems; a more intrinsic, "algorithm-free" analysis will be performed.

In §2 of this paper, the linear algebraic framework of [4] is developed for analytic discrete-time systems, thereby extending the notion of the rank of a system introduced in [10]. This includes the definition of a chain of subspaces constructed from the outputs of the system and their iterates and an analysis of the convergence properties of the chain of subspaces. It is noted that when the function f describing the dynamics is not a submersion, certain new phenomena can occur, requiring a slightly different analysis involving a combination of geometric and algebraic reasoning. Section 3 collects a few results that are useful for establishing relations between some existing work involving rank computations in an "algorithmic" form and the linear algebraic setting proposed here. Section 4 relates the abstract notion of rank, introduced in §2, to the injectivity and surjectivity properties of certain maps strongly connected with left- and right-invertibility. Finally, §5 points out the links between the approach used in this paper and that of [10]; §6 shows the affinity with the work of [28].

2. Rank and structure at infinity. The notion of the *rank* of a nonlinear system was introduced by Fliess in [8] and yielded fundamental results on right- and left-invertibility and noninteracting control of continuous-time systems. Extensions to a class of rational discrete-time systems have been given in [10], using difference algebra in place of differential algebra. Here, using elementary vector space techniques as in [4], the rank of a discrete-time system will be generalized to analytic systems admitting a global state space representation on \mathbb{R}^n . This may be a strong assumption.

2.1. Linear algebraic framework. Consider a discrete-time system

$$(2.1) \quad \Sigma : \begin{aligned}x[k+1] &= f(x[k], u[k]), \\ y[k] &= h(x[k], u[k]),\end{aligned}$$

where $x[k] \in X = \mathbb{R}^n$, $u[k] \in U = \mathbb{R}^m$, $y[k] \in Y = \mathbb{R}^\mu$, f and h are analytic functions of their arguments, and $x[0] = x_0$. It is convenient to let $f^u(x) := f(x, u)$ so that we may write:

$$\begin{aligned}x[1] &= f^{u[0]}(x_0), \\ x[2] &= f^{u[1]}(x[1]) = f^{u[1]} \circ f^{u[0]}(x_0) \\ &\vdots \\ x[k] &= f^{u[k-1]} \circ \dots \circ f^{u[0]}(x_0),\end{aligned}$$

where \circ denotes composition. Then, since $y[k] = h^{u[k]}(x[k])$, where $h^u(x) := h(x, u)$,

$$(2.2) \quad \begin{aligned} y[k] &= y[k](x_0, u[0], \dots, u[k]) \\ &= h^{u[k]} \circ f^{u[k-1]} \circ \dots \circ f^{u[0]}(x_0). \end{aligned}$$

Because (2.1) is time-invariant,

$$(2.3) \quad \begin{aligned} y[k + 1] &= y[k](x[1], u[1], \dots, u[k + 1]) \\ &= y[k](f(x_0, u[0]), u[1], \dots, u[k + 1]), \end{aligned}$$

which will be important later when establishing a certain finiteness property.

Let \mathcal{R}_k denote the ring of real analytic functions of the components of $(x, u[0], \dots, u[k])$, and let \mathcal{K}_k be the associated field of fractions, that is, the field of *meromorphic* functions in the variables $(x, u[0], \dots, u[k])$. A typical element of \mathcal{K}_k would have the form $\eta(v) = \pi(v)/\theta(v)$, where π and θ are elements of \mathcal{R}_k , θ is not the zero function, and $v = (v_1, \dots, v_j)$ denotes the various components of $(x, u[0], \dots, u[k])$. Recall that $\partial/\partial v_i$ acting on η is formally defined by the usual quotient rule of calculus,

$$(2.4) \quad \frac{\partial}{\partial v_i} \frac{\pi(v)}{\theta(v)} := \left(\theta(v) \frac{\partial}{\partial v_i} \pi(v) - \pi(v) \frac{\partial}{\partial v_i} \theta(v) \right) / \theta^2(v),$$

and the formal differential of η is

$$(2.5) \quad d\eta(v) := \sum_{i=1}^j \frac{\partial \eta(v)}{\partial v_i} dv_i.$$

Let \mathcal{E} denote the vector space over $\mathcal{K} := \mathcal{K}_n$ spanned by $\{dx_1, \dots, dx_n, du_1[0], \dots, du_m[0], du_1[n], \dots, du_m[n]\}$. Note that \mathcal{E} is a finite-dimensional vector space; indeed, its dimension is $n + (n + 1)m$. For notational convenience, $\{dx_1, \dots, dx_n\}$ will simply be written as dx , $\{du_1[0], \dots, du_m[0]\}$ as $du[0]$, etc., so that $\mathcal{E} = \text{span}_{\mathcal{K}}\{dx, du[0], \dots, du[n]\}$.

Observe now, for all $0 \leq k \leq n$ and $1 \leq j \leq \mu$, that $dy_j[k] \in \mathcal{E}$, since

$$(2.6) \quad dy_j[k] = \sum_{i=1}^n \frac{\partial y_j[k]}{\partial x_i} dx_i + \sum_{\ell=0}^k \sum_{i=1}^m \frac{\partial y_j[k]}{\partial u_i[\ell]} du_i[\ell].$$

Define a chain of subspaces $\mathcal{E}_0 \subset \dots \subset \mathcal{E}_n$ of \mathcal{E} by [4] (see also [7])

$$(2.7) \quad \mathcal{E}_k := \text{span}_{\mathcal{K}}\{dx, dy[0], \dots, dy[k]\}$$

and the associated list of dimensions $\rho_0 \leq \rho_1 \leq \dots \leq \rho_n$ by

$$(2.8) \quad \rho_k = \dim_{\mathcal{K}} \mathcal{E}_k.$$

We emphasize that $dy[k]$ denotes $\{dy_1[k], \dots, dy_{\mu}[k]\}$ and that this abuse of notation will be used quite often to keep the notation compact.

It will turn out for *generically submersive* systems,¹ that is, for systems where

$$(2.9) \quad n = \text{rank}_{\mathcal{K}} \left[\frac{\partial f}{\partial x}(x, u[0]) : \frac{\partial f}{\partial u}(x, u[0]) \right],$$

¹ The mathematical importance of this assumption will be seen in the next subsection; in terms of control systems, it means that with a feedback, the drift dynamics could be made (generically) invertible, creating a kind of group action.

that

$$(2.10) \quad \rho^* := \rho_n - \rho_{n-1}$$

is a limiting value of the chain (2.7) in the sense that if we were to extend the chain in the obvious manner, then $\rho_{n+r} = \rho_n + r\rho^*$, for all integers $r \geq 0$. Many system models of the form (2.1) would satisfy (2.9), since it is equivalent to $f(\mathbb{R}^n, \mathbb{R}^m)$ having nonempty interior in \mathbb{R}^n , and this is a *necessary* condition for accessibility [17]. It is always satisfied for a time-sampled representation of a continuous-time system. Moreover, it has just been established in [11] that, in a certain sense, rational input-output systems admit *local* state space representations satisfying condition (2.9). To avoid passing to a local representation, the following construction is used here in the general case where (2.9) is not satisfied.

Let $\mathcal{K}^+ := \mathcal{K}_{2n}$, and define $\mathcal{E}^+ := \text{span}_{\mathcal{K}^+} \{dx, du[0], \dots, du[2n]\}$. Define a chain of subspaces $\mathcal{E}_0^+ \subset \dots \subset \mathcal{E}_n^+$ of \mathcal{E}^+ by

$$(2.11) \quad \mathcal{E}_k^+ := \text{span}_{\mathcal{K}^+} \{dx, du[0], \dots, du[n-1], dy[n], \dots, dy[n+k]\}$$

and the associated list of dimensions $\rho_0^+ \leq \dots \leq \rho_n^+$ by

$$(2.12) \quad \rho_k^+ := \dim \mathcal{E}_k^+.$$

Then, even without condition (2.9), it will turn out that

$$(2.13) \quad \rho^{+*} := \rho_n^+ - \rho_{n-1}^+$$

is a limiting value in the sense discussed earlier for ρ^* . Whenever the system (2.1) is generically submersive, it will be established that

$$(2.14) \quad \rho_k^+ = \rho_k + nm, \quad k \geq 0,$$

so that $\rho^{+*} = \rho^*$.

Anticipating these technical results, ρ^{+*} is defined to be the *rank* [10] of the system (2.1).

Remarks. (a) In [6], it is shown (for continuous-time systems) that the chain of subspaces (2.7) is closely related to classical objects in algebra, namely filtrations, and consequently, Hilbert polynomials; a similar result is true in discrete-time [7]. One of the main points of the analysis presented in this paper is the establishment of a priori bounds on the number of steps required to compute the limiting ranks of the filtrations whenever the system has a standard state-space representation; such bounds are not provided by the classical results of algebra, which, on the other hand, apply to more general situations.

(b) In analogy with [4], [22], the list of integers $\{\sigma_0, \dots, \sigma_n\}$ defined by

$$(2.15) \quad \sigma_i = \rho_i - \rho_{i-1}, \quad 0 \leq i \leq n$$

with the convention that $\rho_{-1} = n$, could be called the *transient structure at infinity*, while the list of integers $\{\sigma_0^+, \dots, \sigma_n^+\}$, defined by

$$(2.16) \quad \sigma_i^+ := \rho_i^+ - \rho_{i-1}^+, \quad 0 \leq i \leq n,$$

with the convention that $\rho_{-1}^+ = n + nm$ could be called the *persistent structure at infinity*. For generically submersive systems the two lists coincide, and we can speak simply of the *structure at infinity*; this is also the case for systems satisfying certain constant rank hypotheses in the neighborhood of an equilibrium point, as can be seen from the results of [18], [24], including (constant-coefficient) linear systems.

2.2. Convergence of the chain $\mathcal{E}_0 \subset \dots \subset \mathcal{E}_n$. The goal of this subsection is to justify terminating the chain (2.7) at n , the dimension of the state space of (2.1), whenever the system is generically submersive. For a linear system, this would follow as an easy consequence of the Cayley–Hamilton theorem; in the case of nonlinear systems, more work is required.

Let $k \geq 0$ be any nonnegative finite integer, and recall that \mathcal{K}_k is the field of meromorphic functions of $(x, u[0], \dots, u[k])$. Before, when defining \mathcal{E}_k ($0 \leq k \leq n$), the span was taken with respect to $\mathcal{K} := \mathcal{K}_n$. It is easily seen that the dimension of \mathcal{E}_k does not change if instead the span is taken with respect to \mathcal{K}_k . For $k > n$, define \mathcal{E}_k in the obvious way, following (2.7), taking the span with respect to \mathcal{K}_k .

THEOREM 2.1. *Suppose that (2.1) is generically submersive. Then, for all integers $k \geq n$, $\dim \mathcal{E}_k - \dim \mathcal{E}_{k-1} = \dim \mathcal{E}_n - \dim \mathcal{E}_{n-1}$; that is, $\rho_k - \rho_{k-1} = \rho_n - \rho_{n-1}$.*

The proof of the theorem will be divided into several parts, each establishing a particular property of the chain (2.7) arising from the recursive manner in which the functions $y[k]$ are constructed from the system. Define $\delta : \mathcal{K}_{k-1} \rightarrow \mathcal{K}_k$ by

$$(2.17) \quad (\delta\eta)(x, u[0], \dots, u[k]) = \eta(f(x, u[0]), u[1], \dots, u[k]);$$

it is important that f be generically submersive, for otherwise, $\delta\eta$ may not be a meromorphic function (see (2.20) below). This induces an \mathbb{R} -linear mapping² $\Delta : \text{span}_{\mathcal{K}_{k-1}} \{d\eta | \eta \in \mathcal{K}_{k-1}\} \rightarrow \text{span}_{\mathcal{K}_k} \{d\lambda | \lambda \in \mathcal{K}_k\}$ by

$$(2.18) \quad \begin{aligned} \Delta(d\eta) &:= d(\delta\eta), \\ \Delta(\alpha_1 d\eta_1 + \alpha_2 d\eta_2) &:= \delta(\alpha_1)\Delta(d\eta_1) + \delta(\alpha_2)\Delta(d\eta_2) \end{aligned}$$

for $\eta, \eta_1, \eta_2, \alpha_1, \alpha_2 \in \mathcal{K}_{k-1}$. It should be noted that (2.18) is consistent with the chain rule for differentiation, and that, for instance,

$$(2.19) \quad \begin{aligned} dy_j[k+1] &= d(\delta(y_j[k])) \\ &= \Delta(dy_j[k]) \\ &= \sum_{i=1}^n \delta \left(\frac{\partial y_j[k]}{\partial x_i} \right) dx_i[1] + \sum_{\ell=0}^k \sum_{i=1}^m \delta \left(\frac{\partial y_j[k]}{\partial u_i[\ell]} \right) du_i[\ell+1]. \end{aligned}$$

The following two (equivalent) properties are easily established whenever the system (2.1) is generically submersive:

- P1.** For all $k \geq 1$, $\dim_{\mathcal{K}_{k-1}} \{dx[k]\} = n$.
- P2.** For all $k \geq 0$ and $\forall \eta \in \mathcal{R}_k$, if $\eta \not\equiv 0$, then $\delta(\eta) \not\equiv 0$.

As a consequence, if $\eta \in \mathcal{K}_k$, then $\delta(\eta)$ is well defined. To see the peculiarities a nonsubmersive system may exhibit, consider the example

$$(2.20) \quad \begin{aligned} x[k+1] &= 0, \\ y[k] &= x[k]u[k], \end{aligned}$$

where x, y , and u are in \mathbb{R} . The function $1/x$ is meromorphic, but $\delta(1/x)$ is not defined because $\delta(x) \equiv 0$.

The main ingredients of a proof of Theorem 2.1 are now presented. In the following, if S is a set, then $|S|$ denotes its cardinality.

² The fact that Δ is well defined follows easily from $\Delta(0) = 0$.

LEMMA 2.2. *Suppose that (2.1) is generically submersive and that for some $k \geq 0$, $I_0 \subset I_1 \subset \dots \subset I_k \subset \{1, \dots, \mu\}$ are index sets such that $\mathcal{E}_k = \text{span}_{\mathcal{K}_k} \{dx, dy_{i_0}[0], \dots, dy_{i_k}[k] | i_j \in I_j, 0 \leq j \leq k\}$, and $\dim \mathcal{E}_k = n + |I_0| + \dots + |I_{k-1}| + |I_k|$. Then, $\dim \text{span}_{\mathcal{K}_k} \{dx, dy_{i_0}[0], \dots, dy_{i_k}[k], dy_{i_k}[k+1] | i_j \in I_j, 0 \leq j \leq k\} = n + |I_0| + \dots + |I_{k-1}| + 2|I_k|$. In other words, once an output component becomes independent, it remains independent.*

Proof. For the proof, see Appendix A. □

It immediately follows from the above that there exists a basis of a special form for the chain $\mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots$.

LEMMA 2.3. *Suppose that (2.1) is generically submersive. Then, there exist index sets $I_0 \subset I_1 \subset \dots \subset \{1, \dots, \mu\}$ such that, for all $k \geq 1$, $\{dx, dy_{i_0}[0], \dots, dy_{i_k}[k] | i_j \in I_j, 0 \leq j \leq k\}$ is a basis for \mathcal{E}_k .*

Lemma 2.3 establishes that $\{\sigma_k\} = \{\rho_k - \rho_{k-1}\}$ is a nondecreasing sequence. Since $\sigma_k \leq \min\{m, \mu\}$, it follows that $\{\sigma_k\}$ converges in a finite number of steps. However, this does not allow us to terminate the calculations at the n th step unless the upper bound has already been attained. Considering once again (2.20), we calculate that $(\sigma_0) = 1$, but $(\sigma_1) = 0$; that is, the sequence $\{\sigma_k\}$ of “zeros at infinity of order less than or equal to k ” is not nondecreasing, as is always the case for linear systems and continuous-time nonlinear systems.

LEMMA 2.4. *Suppose that (2.1) is generically submersive, and let I_0, \dots, I_n be as in Lemma 2.3. Then, for each $1 \leq j \leq \mu$, there exists an integer N , $1 \leq N \leq n$, such that*

$$(2.21) \quad dy_j[N] \in \text{span}_{\mathcal{K}_N} \{dy_j[0], \dots, dy_j[N-1], dy_{i_0}[0], \dots, dy_{i_N}[N] | i_k \in I_k, 0 \leq k \leq N\}.$$

Proof. For the proof, see Appendix A. □

The previous and the following lemmas combine to replace the Cayley–Hamilton theorem, which, in the case of a linear system, proves that the chain (2.7) converges in at most n steps.

LEMMA 2.5. *Suppose that (2.1) is generically submersive, and let I_0, \dots, I_n be as in Lemma 2.3. Suppose that $1 \leq j \leq \mu$, and let N be as in Lemma 2.4. Then, for all $k \geq N$,*

$$(2.22) \quad dy_j[k] \in \text{span}_{\mathcal{K}_k} \{dy_j[0], \dots, dy_j[N-1], dy_{i_0}[0], \dots, dy_{i_k}[k]\} \\ \text{for } 1 \leq s \leq N, i_s \in I_s, \text{ and for } s > N, i_s \in I_N\}.$$

Proof. The proof is immediate from (2.18) and Lemma 2.4. □

The proof of Theorem 2.1 is now given easily. Let $\{I_k\}$ be the collection of index sets determined by Lemma 2.2. By Lemma 2.5, $I_{n+r} = I_n$ for all $r \geq 1$. Hence, the components of the output either become independent by the n th iteration of the dynamics, or they remain dependent for all iterations. Consequently, for all $r \geq 1$, $\dim \mathcal{E}_{n+r} - \dim \mathcal{E}_{n+r-1} = \dim \mathcal{E}_n - \dim \mathcal{E}_{n-1} = \rho^*$.

2.3. Convergence of the chain $\mathcal{E}_0^+ \subset \mathcal{E}_1^+ \subset \dots$. This section addresses the convergence properties of the chain $\mathcal{E}_0^+ \subset \mathcal{E}_1^+ \subset \dots$ for systems that are not necessarily submersive. The idea behind the analysis is that the effect of the nilpotent part of the system on the output sequence is short lived and can be eliminated from the analysis by “ignoring” the first n time instances of the output; this is essentially what

is accomplished by including $\{du[0], \dots, du[n-1]\}$ in the definition of $\mathcal{E}_0^+ \subset \mathcal{E}_1^+ \subset \dots$. The analytical aspects of the proof are based on the following construction.

Let $M_0 := \mathbb{R}^n$ and define for $k \geq 1$

$$(2.23) \quad M_k := x[k] (\mathbb{R}^n \times (\mathbb{R}^m)^k).$$

Observe that $M_0 \supset M_1 \supset \dots$ since

$$(2.24) \quad M_{k+1} = f(M_k, \mathbb{R}^m).$$

Define $d_0 := n$ and for $k \geq 1$,

$$(2.25) \quad d_k := \text{rank}_{\mathcal{K}_{k-1}} x[k] := \dim \text{span}_{\mathcal{K}_{k-1}} \{dx[k]\}.$$

LEMMA 2.6. *The sequence of integers $\{d_k\}$ is nonincreasing, and if $d_k = d_{k-1}$ then $d_{k+1} = d_k$. Consequently, since d_j can decrease at most n -times, $d_j = d_n$ for all $j \geq n$.*

The proof is given in Appendix B. Very roughly speaking, Lemma 2.6 says that “ $f : M_k \times \mathbb{R}^m \rightarrow M_{k+1}$ behaves like a ‘submersion’ for $k \geq n$ since the ‘dimension’ of M_k is d_k .” Of course, M_k , in general, does not have the structure of a manifold, hence, the imprecision of such a statement. Nevertheless, if we pursue this line of thought for a moment and suppose that $M_0 \supset M_1 \supset \dots$ is a nested sequence of embedded analytic submanifolds, it is clear that, if $\dim M_k = \dim M_{k-1}$, then $\dim M_{k+1} = \dim M_k$, because the condition $\dim M_k = \dim M_{k-1}$ implies that $f : M_{k-1} \times \mathbb{R}^m \rightarrow M_k$ is a submersion. This combined with M_k being open in M_{k-1} gives the result. The proof in Appendix B makes this line of reasoning rigorous with a local analysis, which also establishes the following result: For $k \geq 2$, $j \geq 0$ define $\delta^k : \mathcal{R}_j \rightarrow \mathcal{R}_{j+k}$ by, if $\alpha \in \mathcal{R}_j$, and $0 \leq i \leq k-1$

$$(2.26) \quad \delta^{i+1}(\alpha) = \delta(\delta^i(\alpha)).$$

LEMMA 2.7. *Let $k \geq 0$, $\alpha \in \mathcal{R}_k$. If $\delta^n(\alpha) \neq 0$ (i.e., is not the identically zero function), then $\delta^{n+j}(\alpha) \neq 0$ for all $j \geq 1$.*

Let $\mathcal{R}_k^{\delta^n}$ be the set of real analytic functions

$$(2.27) \quad \mathcal{R}_k^{\delta^n} := \{\delta^n(\alpha) | \alpha \in \mathcal{R}_k\}.$$

$\mathcal{R}_k^{\delta^n} \subset \mathcal{R}_{n+k}$ as a subring, and thus, the associated set of fractions, denoted $\mathcal{K}_k^{\delta^n}$, is a field; indeed, it is a subfield of \mathcal{K}_{n+k} . Note that $\mathcal{K}_k^{\delta^{n+1}} \subset \mathcal{K}_{k+1}^{\delta^n}$.

By Lemma 2.7, for each $k \geq 0$, the two mappings

$$(2.28) \quad \delta : \mathcal{K}_k^{\delta^n} \rightarrow \mathcal{K}_k^{\delta^{n+1}} \subset \mathcal{K}_{k+1}^{\delta^n}$$

and

$$(2.29) \quad \Delta : \text{span}_{\mathcal{K}_k^{\delta^n}} \{d\lambda | \lambda \in \mathcal{K}_k^{\delta^n}\} \rightarrow \text{span}_{\mathcal{K}_{k+1}^{\delta^{n+1}}} \{d\eta | \eta \in \mathcal{K}_k^{\delta^{n+1}}\} \subset \text{span}_{\mathcal{K}_{k+1}^{\delta^n}} \{d\gamma | \gamma \in \mathcal{K}_{k+1}^{\delta^n}\}$$

can be defined as in (2.17) and (2.18), respectively.

The final step of the analysis is to reduce the study of the chain $\mathcal{E}_0^+ \subset \mathcal{E}_1^+ \subset \dots$ to that of a related chain to which the proof technique of §2.2, that is, Lemmas 2.2–2.5,

can be applied with only minor modifications. As in §2.2, it is necessary to slightly modify the definition of \mathcal{E}_k^+ without changing its dimension:

$$(2.30) \quad \mathcal{E}_k^+ := \text{span}_{\mathcal{K}_{n+k}} \{dx, du[0], \dots, du[n-1], dy[n], \dots, dy[n+k]\},$$

for all $k \geq 0$, and

$$(2.31) \quad \mathcal{E}_{-1}^+ := \text{span}_{\mathcal{K}_{n-1}} \{dx, du[0], \dots, du[n-1]\}.$$

Note that $\{dx[n]\} \subset \mathcal{E}_{-1}^+$. Let $L \subset \{1, \dots, n\}$ be such that $\{dx_i[n] | i \in L\}$ is a basis for $\text{span}_{\mathcal{K}_{n-1}} \{dx[n]\}$, and let \mathcal{W} be such that

$$(2.32) \quad \mathcal{E}_{-1}^+ = \text{span}\{dx_i[n] | i \in L\} \oplus \mathcal{W}.$$

Introduce

$$(2.33) \quad \begin{aligned} \tilde{\mathcal{S}}_k &:= \text{span}_{\mathcal{K}_{n+k}} \{dx_i[n], dy[n], \dots, dy[n+k] | i \in L\}, \\ \tilde{\mathcal{S}}_{-1} &:= \text{span}_{\mathcal{K}_{n-1}} \{dx_i[n] | i \in L\}. \end{aligned}$$

Since $y[n+j](x, u[0], \dots, u[n+j]) = y[j](x[n], u[n], \dots, u[n+j])$ for $j \geq 0$, it follows that

$$(2.34) \quad \mathcal{E}_k^+ = \tilde{\mathcal{S}}_k \oplus \mathcal{W}, \quad k \geq 1.$$

Hence, for $k \geq 0$,

$$(2.35) \quad \sigma_k^+ = \dim \tilde{\mathcal{S}}_k - \dim \tilde{\mathcal{S}}_{k-1}.$$

The reason for doing all of this is that the generators of $\tilde{\mathcal{S}}_k$, for $k \geq 1$, are elements of $\{d\lambda | \lambda \in \mathcal{K}_k^{\delta^n}\}$. Hence, letting

$$(2.36) \quad \begin{aligned} \mathcal{S}_k &:= \text{span}_{\mathcal{K}_k^{\delta^n}} \{dx_i[n], dy[n], \dots, dy[n+k] | i \in L\}, \\ \mathcal{S}_{-1} &:= \text{span}_{\mathcal{K}_{n-1}} \{dx_i[n] | i \in L\}, \end{aligned}$$

it follows that, for $k \geq 0$,

$$(2.37) \quad \sigma_k^+ = \dim \mathcal{S}_k - \dim \mathcal{S}_{k-1}.$$

Moreover, even though $\mathcal{S}_k \subset \mathcal{S}_{k+1}$ is *not* a subspace of \mathcal{S}_{k+1} ,

$$(2.38) \quad \dim \text{span}_{\mathcal{K}_{k+1}^{\delta^n}} \{\mathcal{S}_k\} = \dim \mathcal{S}_k,$$

and therefore, the proofs of the obvious modifications of Lemmas 2.2–2.5 for the chain $\mathcal{S}_0 \subset \mathcal{S}_1 \subset \dots$ go through with only minor changes, which will not be repeated here.

THEOREM 2.8. *For all integers $k \geq n$, $\dim \mathcal{E}_k^+ - \dim \mathcal{E}_{k-1}^+ = \dim \mathcal{E}_n^+ - \dim \mathcal{E}_{n-1}^+$; that is, $\rho_k^+ - \rho_{k-1}^+ = \rho_n^+ - \rho_{n-1}^+$. Moreover, whenever (2.1) is generically submersive, the ordered lists $\{\sigma_0, \sigma_1, \dots, \sigma_n\}$ and $\{\sigma_0^+, \dots, \sigma_n^+\}$ are equal.*

The last part of the theorem follows from the fact that when (2.1) is generically submersive, the index set L in (2.32) is equal to $\{1, \dots, n\}$. Then, since $y[n+j](x, u[0], \dots, u[n+j])$ can be expressed as $y[j](x[n], u[n], \dots, u[n+j])$, \mathcal{E}_k and \mathcal{S}_k are naturally isomorphic under $x \rightarrow x[n], u[0] \rightarrow u[n], \dots, u[k] \rightarrow u[n+k]$.

3. Further characterizations of the rank and structure at infinity. This section and the rest of the paper will concentrate on generically submersive systems. Similar results, as per the development of §2.3, can be stated for the general case.

3.1. Jacobian matrices. The goal here is to provide a computationally convenient means of evaluating the rank ρ^* . The same result is also useful for showing the invariance of ρ^* under the action of invertible (static or dynamic) state variable feedback.

Following [14], which, in turn, was based upon [23], consider the Jacobian matrices

$$(3.1) \quad J_k(x, u[0], \dots, u[k]) := \frac{\partial(y[0], \dots, y[k])}{\partial(u[0], \dots, u[k])},$$

for $0 \leq k \leq n$, and their associated ranks

$$(3.2) \quad R_k := \text{rank}_{\mathcal{K}} J_k.$$

Note that the matrices J_k can be evaluated symbolically; their ranks can be evaluated numerically since the rank over \mathcal{K} is the same as the generic rank considered in [23].

Applying arguments identical to those used in [4, §2.1] results in the following relation between the integers ρ_k and R_k .

PROPOSITION 3.1. *For each $0 \leq k \leq n$, $\rho_k = n + R_k$. Hence, if (2.1) is generically submersive, then $\rho^* = R_n - R_{n-1}$.*

A quite different way of obtaining a result similar to the first part of Proposition 3.1 is given in [6].

Consider now a discrete-time linear system

$$(3.3) \quad \begin{aligned} x[k+1] &= Ax[k] + Bu[k], \\ y[k] &= Cx[k] + Du[k]. \end{aligned}$$

Then the Jacobian matrix J_k is given by the usual Toeplitz matrix

$$(3.4) \quad J_k = \begin{bmatrix} D & 0 & 0 \\ CB & D & \\ CAB & CB & \vdots \\ \vdots & & 0 \\ CA^{k-1}B & CA^{k-2}B & \dots & D \end{bmatrix}.$$

The results of [21] and [26] in conjunction with Proposition 3.1 justify the terminology adopted in §2.2 concerning the rank and structure at infinity of a nonlinear system.

The following is the analogue of [10, III.B.2. Proposition].

COROLLARY 3.2. *In the case of a linear system, the rank ρ^* defined by (2.10) agrees with the classical rank of the transfer matrix. Moreover, the list of integers $\{\sigma_0, \dots, \sigma_n\}$ defined in (2.15) is precisely the structure at infinity as it is normally defined on the basis of the transfer matrix [21], [26].*

REMARK 3.3. For a linear system, it is easy to verify that the lists $\{\sigma_k^+\}$ and $\{\sigma_k\}$ coincide, whether or not the system is generically submersive.

3.2. A related chain of subspaces. Related to the chain $\mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots$ is the chain $\mathcal{H}_0 \subset \mathcal{H}_1 \subset \dots$ defined solely in terms of the output [4]:

$$(3.5) \quad \mathcal{H}_k = \text{span}_{\mathcal{K}_k} \{dy[0], \dots, dy[k]\}.$$

It also can be used to determine the rank of the system, and this will be important for making contact with the fundamental work of [10].

THEOREM 3.4. *Suppose that (2.1) is generically submersive. For all integers $k \geq n$, $\rho^* = \dim \mathcal{H}_k - \dim \mathcal{H}_{k-1}$.*

Proof. By Lemma 2.5, for $k \geq n$ $\rho^* \geq \dim \mathcal{H}_k - \dim \mathcal{H}_{k-1}$. On the other hand, for $1 \leq j$, $\mathcal{H}_j = \mathcal{H}_{j-1} + \text{span}\{dy[j]\}$, $\mathcal{E}_j = \mathcal{E}_{j-1} + \text{span}\{dy[j]\}$ and $H_0 \subset \mathcal{E}_0$. Thus, $\dim \mathcal{H}_j - \dim \mathcal{H}_{j-1} \geq \dim \mathcal{E}_j - \dim \mathcal{E}_{j-1}$. \square

3.3. Remarks on the inversion algorithm. The importance of the inversion algorithm of Singh [27], which is an extension to nonlinear continuous-time systems of the well-known algorithm of Silverman [26], need not be underlined here. The algorithm has also been used in the study of discrete-time nonlinear systems [18] and [19], but always expressed in a form involving the implicit function theorem. Consequently, the results of the algorithm can be difficult to interpret unless one remains in a neighborhood of an equilibrium point. This problem can be removed by working at the level of the differentials of the outputs, which linearizes the computations and allows the analysis of [4] to be carried through to the discrete-time setting. Since the algorithm in the form we will use it has already appeared in several publications for continuous-time systems [3], [4], [16], the basic idea will only be sketched here by giving the first steps of the algorithm. Establishing the validity and convergence properties of the algorithm is quite easy using the analysis of §2.2.

It is assumed that (2.1) is generically submersive; an extension to general systems can be envisioned along the lines of §2.3.

Step 0. Calculate $dy[0]$ and write it as

$$(3.6) \quad dy[0] = a_0(x, u[0])dx + b_0(x, u[0])du[0] .$$

Define

$$(3.7) \quad s_0 := \text{rank}_{\mathcal{K}_0} b_0 .$$

Permute, if necessary, the components of y so that the first s_0 rows of b_0 are linearly independent. Decompose y so that

$$(3.8) \quad dy[0] = \begin{bmatrix} d\tilde{y}_0[0] \\ d\hat{y}_0[0] \end{bmatrix} = \begin{bmatrix} \tilde{a}_0 \\ \hat{a}_0 \end{bmatrix} dx + \begin{bmatrix} \tilde{b}_0 \\ \hat{b}_0 \end{bmatrix} du[0],$$

where \tilde{y}_0 has s_0 rows. Since the rows of \hat{b}_0 are \mathcal{K}_0 -dependent on the rows of \tilde{b}_0 , there exists a matrix $M_0(x, u[0])$ with entries in \mathcal{K}_0 such that

$$(3.9) \quad \hat{b}_0 = M\tilde{b}_0,$$

and thus,

$$(3.10) \quad \begin{aligned} d\hat{y}_0[0] &= \hat{a}_0 dx + M_0\{d\tilde{y}_0[0] - \tilde{a}_0 dx\} \\ &=: \bar{a}_0 dx + \bar{b}_0 d\tilde{y}_0[0]. \end{aligned}$$

End of Step 0.

Step 1. Compute

$$(3.11) \quad \begin{aligned} d\hat{y}_0[1] &= (\delta\bar{a}_0) dx[1] + (\delta\bar{b}_0) d\tilde{y}_0[1] \\ &=: a_1(x, u[0], u[1]) dx + b_1(x, u[0], u[1]) du[0] + c_1(x, u[0], u[1]) d\tilde{y}_0[1]. \end{aligned}$$

Define

$$(3.12) \quad s_1 := \text{rank}_{\mathcal{K}_1} \begin{bmatrix} \tilde{b}_0 \\ b_1 \end{bmatrix}$$

and repeat the basic operations of Step 1; see [3], [4], and [16], for example.

The validation of the steps in the algorithm is achieved by noting that it produces a basis for $\mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots$, and thus, by Lemma 2.2, since $\{dx, d\tilde{y}_0[0]\}$ is a linearly independent set, so is $\{dx, d\tilde{y}_0[0], d\tilde{y}_0[1]\}$, etc. Its convergence in no more than n steps follows from Theorem 2.1.

A similar connection with the interesting work of [24] on dynamic feedback solutions to the noninteracting control problem could be pursued also along the lines already clearly established in [4].

4. Invertibility. A linear system is usually said to be right-invertible if the rank of its transfer matrix is equal to the number of output components, and left-invertible if its rank equals the number of input components. Systemically, right-invertibility means that by a proper choice of the initial condition *and* input sequence, any output sequence can be generated; that is, the map from initial conditions and inputs is onto $Y^\infty = Y \times Y \times \dots$, the space of all output sequences. Left-invertibility is equivalent to injectivity of the map from inputs to outputs, for a fixed initial condition.

In the case of nonlinear systems, though such global notions of invertibility are attractive, simple examples show the difficulty of trying to say anything intelligent about them; hence, we are led to localizing the concepts. Following [3], for $k \geq 0$, let $H_k : X \times U^{k+1} \rightarrow Y^{k+1}$ be the map that sends $(x, u[0], \dots, u[k])$ to $(y[0], \dots, y[k])$, and let $E_k : X \times U^{k+1} \rightarrow X \times Y^{k+1}$ by $(x, u[0], \dots, u[k]) \mapsto (x, y[0], \dots, y[k])$.

DEFINITION 4.1. The system (2.1) is *almost everywhere locally surjective*, if, for every $k \geq 0$, the image of H_k has nonempty interior. The system is *almost everywhere locally injective*, if, for every $k \geq 0$, there exists an open and dense subset \mathcal{O}_k of $X \times U^{k+1}$ with the property that, for each point $p = (x, u[0], \dots, u[k]) \in \mathcal{O}_k$ there exists an open neighborhood of p , $\tilde{\mathcal{O}}_k(p)$, and an analytic insertion³ $i_k : \tilde{\mathcal{O}}_k(p) \rightarrow X \times U^{k+1} \times U^n$ such that if $p_1, p_2 \in \tilde{\mathcal{O}}_k(p)$ and $E_{n+k}(i_k(p_1)) = E_{n+k}(i_k(p_2))$, then $p_1 = p_2$.

These properties can be characterized as follows.

THEOREM 4.2. *Assume that the nonlinear system (2.1) is generically submersive. Then the system is almost everywhere locally surjective if and only if any one of the following equivalent conditions is satisfied:*

- (a) for all $k \geq 0$, $\dim \text{span}\{dy[0], \dots, dy[k]\} = (k + 1)\mu$;
- (b) $\dim \text{span}\{dy[0], \dots, dy[n]\} = (n + 1)\mu$;
- (c) $\rho^* = \mu$; i.e., the rank of the system equals the number of output components.

The system is almost everywhere locally injective if, and only if, any one of the following equivalent conditions is satisfied:

- (d) for all $k \geq 0$, $\{du[0], \dots, du[k]\} \subset \text{span}\{dx, dy[0], \dots, dy[k + n]\}$
- (e) $du[0] \subset \text{span}\{dx, dy[0], \dots, dy[n]\}$;
- (f) $\rho^* = m$; i.e., the rank of the system equals the number of input components.

Proof. (a) \implies (b) is immediate. (b) \implies (c) is given by Theorem 3.4. (c) \implies (a) follows from the same kind of reasoning employed in proving Lemmas 2.4 and 2.5 and is not repeated here. It suffices to show that (a) is equivalent to almost everywhere

³ That is, if τ_k represents that natural projection of $X \times U^{k+1} \times U^n$ onto $X \times U^{k+1}$, then $\tau_k \circ i_k|_{\tilde{\mathcal{O}}_k(p)}$ is the identity.

local surjectivity. The image of H_k has nonempty interior if, and only if, there exists an open set of points where the rank of H_k over the reals equals $(k + 1)\mu$. This is equivalent to H_k having rank $(k + 1)\mu$ over \mathcal{K}_k , which is equivalent to (a). Turning to almost everywhere local injectivity, (d) \implies (e) is evident. (e) \iff (f) is Corollary A.2 in Appendix A. It is now shown that (e) \implies (d): Applying Δ to both sides of (e) yields

$$du[1] \subset \text{span}\{dx[1], dy[1], \dots, dy[n + 1]\} .$$

Hence, adding $\text{span}\{dx, du[0], dy[0]\}$ to the right-hand side,

$$\begin{aligned} du[1] &\subset \{dx, du[0], dy[0], dy[1], \dots, dy[n + 1]\} \\ &\subset \text{span}\{dx, dy[0], \dots, dy[n + 1]\}, \end{aligned}$$

where (e) has been used in the last step. The remainder of a proof by induction is clear. To finish up, it suffices now to show that (d) is equivalent to almost everywhere local injectivity. Without loss of generality, it can be assumed that the neighborhood \tilde{O}_k is such that $E_{n+k} \circ i_k : \tilde{O}_k \rightarrow X \times Y^{n+k+1}$ has constant rank. Then almost everywhere local injectivity means that this rank equals $n + (k + 1) \cdot m$. This is equivalent to $\text{rank}_{\mathcal{K}_{n+k}} (\partial(y[0], \dots, y[n + k])/\partial(u[0], \dots, u[k])) = (k + 1)m$, which is equivalent to (d). \square

Remark. A quite different approach to invertibility is taken in [10]; the results of the next section show that, in the case of polynomial systems, the two approaches coincide.

5. Difference algebra and the transformal transcendence degree. The purpose of this section is to prove that the rank ρ^* defined in (2.10), when specialized to systems whose right-hand side depends polynomially on x and u (more precisely, on their components), corresponds to the transformal transcendence degree used in [10].

Consider a system

$$(5.1) \quad \begin{aligned} \Sigma_{P,Q} \quad x[k + 1] &= P(x[k], u[k]) \\ y[k] &= Q(x[k], u[k]), \end{aligned}$$

where $P : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$ and $Q : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^\mu$ are such that each of their components is a polynomial of x and u , with coefficients in \mathbb{R} . This system is clearly analytic, so the analysis of §2 applies. For the purpose of clarity in presenting the results, it will be assumed that (5.1) is generically submersive. An extension to rational systems could also be undertaken.

The following definition, adapted from [10], should actually be *derived* by constructing the difference field associated to $\Sigma_{P,Q}$ and then applying the definition used in [10].

DEFINITION 5.1. The transformal transcendence degree of $\Sigma_{P,Q}$, denoted $d^0(\Sigma_{P,Q})$ equals the maximal number of components of y , say $\{y_{i_1}, \dots, y_{i_\rho}\}$, such that for any $k \geq 0$ there does not exist any nontrivial polynomial π with coefficients in \mathbb{R} such that

$$\pi(y_{i_1}[0], \dots, y_{i_\rho}[0], \dots, y_{i_1}[k], \dots, y_{i_\rho}[k]) = 0.$$

In other words, for any $k \geq 0$, $y_{i_1}[0], \dots, y_{i_\rho}[0], \dots, y_{i_1}[k], \dots, y_{i_\rho}[k]$ viewed as polynomials of $x, u[0], \dots, u[k]$ are algebraically independent.

For those readers familiar with Kähler differentials, the equality $\rho^* = d^0(\Sigma_{P,Q})$ is immediate from Theorem 3.4. For the benefit of other readers, an independent, straightforward proof is given. Lemma 5.2, which follows, is well-known, though hard to find in the form presented (cf. [28]).

Let $v = (v_1, \dots, v_r)$ be an r -tuple of indeterminants, let $\mathbb{R}[v]$ denote the ring of polynomials of (v_1, \dots, v_r) with coefficients in \mathbb{R} and let $\mathbb{R}(v)$ be the corresponding field of rational functions. Define a vector space over $\mathbb{R}(v)$ by $V := \text{span}\{dv_1, \dots, dv_r\}$ and define the mapping $d : \mathbb{R}(v) \rightarrow V$ by

$$(5.2) \quad d\left(\frac{p(v)}{q(v)}\right) := \frac{1}{(q(v))^2} \sum_{j=1}^r \left(\frac{\partial p(v)}{\partial v_j} q(v) - p(v) \frac{\partial q(v)}{\partial v_j} \right) dv_j$$

in the usual way; see [30, Chap. 5, §10.5] for how to define a differential calculus of rational functions without taking limits.

LEMMA 5.2. *A collection of polynomials $\{P_1, \dots, P_k\} \subset \mathbb{R}[v]$ is algebraically independent if, and only if, the set $\{dP_1, \dots, dP_k\}$ is linearly independent in $(V, \mathbb{R}(v))$.*

Proof. Suppose that $\{P_1, \dots, P_k\}$ is algebraically independent; then $k \leq r$. Assume first that $k = r$. $\{P_1, \dots, P_r\}$ is then a basis for $\mathbb{R}[v]$, and consequently, for each $1 \leq i \leq r$, there is a nontrivial polynomial $Q_i(\lambda_1, \dots, \lambda_{r+1})$ such that $Q_i(v_i, P_1, \dots, P_r) = 0$.

PROPOSITION 5.3. *For each $1 \leq i \leq r$, the polynomial $\partial Q_i / \partial \lambda_1$ is nontrivial.*

Proof. Suppose it is trivial. Then $Q_i(v_i, P_1, \dots, P_r) = Q_i(0, P_1, \dots, P_r) =: \bar{Q}_i(P_1, \dots, P_r)$. Thus, \bar{Q}_i must be a trivial polynomial because $\{P_1, \dots, P_r\}$ is algebraically independent. It follows that Q_i is a trivial polynomial, which contradicts its definition. \square

Continuing with the proof of Lemma 5.2, since $0 = Q_i(v_i, P_1, \dots, P_r)$,

$$(5.3) \quad 0 = d(Q_i(v_i, P_1, \dots, P_r)) = \frac{\partial Q_i}{\partial \lambda_1}(v_i, P_1, \dots, P_r) dv_i + \sum_{j=1}^r \frac{\partial Q_i}{\partial \lambda_{j+1}}(v_i, P_1, \dots, P_r) dP_j;$$

see [30, Chap. 5] for the chain rule. By Proposition 5.3, $\partial Q_i / \partial \lambda_1$ is nontrivial, and thus,

$$(5.4) \quad dv_i = \sum_{j=1}^r k_{ij} dP_j,$$

where

$$k_{ij} := \left(\frac{\partial Q_i}{\partial \lambda_1}(v_i, P_1, \dots, P_r) \right)^{-1} \left(\frac{\partial Q_i}{\partial \lambda_{j+1}}(v_i, P_1, \dots, P_r) \right) \in \mathbb{R}(v).$$

Thus, $\text{span}\{dv_1, \dots, dv_r\} \subset \text{span}\{dP_1, \dots, dP_r\}$, proving the linear independence of $\{dP_1, \dots, dP_r\}$.

If $k < r$, then there exist P_{k+1}, \dots, P_r such that $\{P_1, \dots, P_r\}$ is a basis for $\mathbb{R}(v)$. From the above, $\{dP_1, \dots, dP_r\}$ is linearly independent, and therefore, so must be $\{dP_1, \dots, dP_k\}$.

To prove the other direction of the lemma, suppose that $\{P_1, \dots, P_k\}$ is algebraically dependent. Then there exists a nontrivial polynomial $Q(\lambda_1, \dots, \lambda_k)$ with coefficients in \mathbb{R} such that $Q(P_1, \dots, P_k) = 0$. Hence, $0 = \sum_{j=1}^k \partial Q / \partial \lambda_j (P_1, \dots, P_k) dP_j$,

proving that $\{dP_1, \dots, dP_k\}$ is a linearly dependent set in $(V, \mathcal{R}(v))$. This completes the proof of Lemma 5.2. \square

The constructions and results of §2 hold clearly for polynomial systems (5.1) with the field \mathcal{K} replaced⁴ by $\mathcal{R}[x, u[0], \dots, u[n]]$. This observation, combined with Lemma 5.2 and Theorem 3.4, yields the following result.

THEOREM 5.4. *For the polynomial system (5.1), $\rho^* = d^0(\Sigma_{P,Q})$.*

Remark. Lemma 5.2 can be equivalently stated as: The following two conditions are equivalent:

(a) There exists a nontrivial polynomial π such that $\pi(P_1, \dots, P_r) = 0$

(b) The set $\{dP_1, \dots, dP_r\}$ is linearly dependent in $(V, \mathcal{R}(v))$

The implication (a) \implies (b) remains true with π, P_1, \dots, P_r replaced by analytic functions. However, the converse is then true only *locally*, and even then only on subsets where certain constant dimensional conditions are met; indeed, this is the well-known Rank Theorem.

6. Relation to generic observation fields. The chain of subspaces $\mathcal{E}_0 \subset \dots \subset \mathcal{E}_k \subset \dots$ is measuring how the input components are appearing in the outputs, assuming that the initial state is known. In a similar manner, one could study how the initial state components appear in the outputs, assuming that the inputs are known. This is called *observability* [15], and in the context of the formalism of this paper, could be studied via the chain $\mathcal{O}_0 \subset \dots \subset \mathcal{O}_k \subset \dots$, where

$$(6.1) \quad \mathcal{O}_k := \text{span}_{\mathcal{K}_k} \{du[0], \dots, du[k], dy[0], \dots, dy[k]\}.$$

Then, as in [4] and this paper, a connection could be established between the ranks of certain Jacobian matrices, the dimensions of the subspaces \mathcal{O}_k and/or the transcendence degree of a certain (differential) field. This (plus a whole lot more, such as the construction of a realization theory for polynomial input-output maps) was done by Sontag [28] in 1979 for a very general class of discrete-time polynomial systems (cf. his generic observation fields, Q_f^K). An extension to analytic systems would follow along the lines of [31]. More recent work on the analysis of the observability of continuous-time systems by algebraic means can be found in [5] and the references therein.

Appendix A.

Proof of Lemma 2.2. For the sake of the study of invertibility in §3, it is useful to prove a little more than is required by the lemma. The following notation is used only in Appendix A; it will help to keep the formulas concise. If M is a subset of \mathcal{E} , then $[M]$ denotes its *span* (see [20, p. 16]). If $\{v_1, \dots, v_s\}$ is a set of *linearly independent* elements in \mathcal{E} , this will be denoted by $\{v_1, \dots, v_s\}^*$. For example, let $I_0 \subset \{1, \dots, \mu\}$ be such that $\{dx, dy_{i_0}[0] | i_0 \in I_0\}$ is a basis for \mathcal{E}_0 . Then, this can be succinctly stated as $[\{dx, dy_{i_0}[0] | i_0 \in I_0\}^*] = \mathcal{E}_0$.

Suppose that for some $0 \leq k$, index sets $I_0 \subset I_1 \subset \dots \subset I_k \subset \{1, \dots, \mu\}$ and $\{1, \dots, m\} \supset J_0 \supset J_1 \supset \dots \supset J_k$ have been selected so that for each $0 \leq t \leq k$,

$$(A.1) \quad \mathcal{E}_t = [\{dx, dy_{i_r}[r] | 0 \leq r \leq t, i_r \in I_r\}^*]$$

$$(A.2) \quad [dx, du[0], \dots, du[t]] = [\{dx, dy_{i_r}[r], du_{j_{t-r}}[r] | 0 \leq r \leq t, i_r \in I_r, j_r \in J_r\}^*].$$

⁴ So that the proof of Lemma 2.4, which used the Rank Theorem, can be carried over, one must note that (2.21) holds for (5.1) over \mathcal{K} if, and only if, it holds over $\mathcal{R}[x, u[0], \dots, u[n]]$.

and

$$(A.3) \quad du[0] \subset \mathcal{E}_t \oplus \{ \{ du_j[0] | j \in J_t \} \}.$$

That this is possible for $k = 0$ is obvious. It will first be shown that $I_{k+1} \supset I_k$ can be chosen so that (A.1) holds for $0 \leq t \leq k + 1$.

CLAIM A.1. *The set*

$$\{ dx, dy_{i_r}[r], dy_{i_k}[k + 1], du_{j_k}[0], du_{j_{k-r}}[r + 1] | 0 \leq r \leq k, i_r \in I_r, j_r \in J_r \}$$

is linearly independent.

Proof. To see how the arguments go, consider first (A.2) for $t = 0$ and apply Δ to both sides to obtain

$$[dx[1], du[1]] = [\{ dx[1], dy_{i_0}[1], du_{j_0}[1] | i_0 \in I_0, j_0 \in J_0 \}].$$

Adding $[dx, du[0]]$ to both sides yields

$$[dx, du[0], du[1]] = [\{ dx, du[0], dy_{i_0}[1], du_{j_0}[1] | i_0 \in I_0, j_0 \in J_0 \}].$$

Applying (A.2) for $t = 0$ results in

$$[dx, du[0], du[1]] = [\{ dx, dy_{i_0}[0], du_{j_0}[0], dy_{i_0}[1], du_{j_0}[1] | i_0 \in I_0, j_0 \in J_0 \}].$$

The independence of the vectors on the left-hand side implies the independence of those on the right-hand side by counting the number of elements. In particular, the vectors $\{ dx, dy_{i_0}[0], dy_{i_0}[1], | i_0 \in I_0, \}$ are linearly independent, and thus, one can choose $I_1 \supset I_0$ such that (A.1) holds for $0 \leq t \leq 1$.

In general, consider (A.2) for $t = k$ and apply Δ to both sides to obtain

$$(A.4) \quad [dx[1], du[1], \dots, du[k + 1]] \\ = [\{ dx[1], dy_{i_r}[r + 1], du_{j_{k-r}}[r + 1] | 0 \leq r \leq k, i_r \in I_r, j_r \in J_r \}].$$

Add $[dx, du[0]]$ to both sides to obtain

$$[dx, du[0], \dots, du[k + 1]] \\ = [\{ dx, du[0], dy_{i_r}[r + 1], du_{j_{k-r}}[r + 1] | 0 \leq r \leq k, i_r \in I_r, j_r \in J_r \}^*].$$

The independence of the vectors on the left-hand side implies the independence of those on the right-hand side by counting the number of elements. Applying (A.2) first for $t = 0$, and then successively for $t = 1, \dots, t = k$ results in

$$(A.5) \quad [dx, du[0], \dots, du[k + 1]] \\ = [\{ dx, dy_{i_0}[0], du_{j_0}[0], dy_{i_r}[r + 1], du_{j_{k-r}}[r + 1] | 0 \leq r \leq k, i_r \in I_r, j_r \in J_r \}^*] \\ = [\{ dx, dy_{i_0}[0], dy_{i_1}[1], du_{j_1}[0], dy_{i_s}[s + 1], du_{j_{k-r}}[r + 1] | 0 \\ \leq r \leq k, 1 \leq s \leq k, i_s \in I_s, j_r \in J_r \}^*] \\ = [\{ dx, dy_{i_0}[0], dy_{i_1}[1], dy_{i_2}[2], du_{j_2}[0], dy_{i_s}[s + 1], du_{j_{k-r}}[r + 1] | 0 \\ \leq r \leq k, 2 \leq s \leq k, i_r \in I_r, j_r \in J_r \}^*] \\ = \text{for each } 0 \leq t \leq k \\ = [\{ dx, dy_{i_0}[0], \dots, dy_{i_t}[t], du_{j_t}[0], dy_{i_s}[s + 1], du_{j_{k-r}}[r + 1] | 0 \\ \leq r \leq k, t \leq s \leq k, i_r \in I_r, j_r \in J_r \}^*]. \quad \square$$

Hence, we can choose $I_{k+1} \supset I_k$ so that (A.1) holds for each $0 \leq t \leq k + 1$. It is next shown that $J_{k+1} \subset J_k$ can be chosen such that (A.2) and (A.3) hold. From (A.3) for $t = k$, since $\mathcal{E}_k \subset \mathcal{E}_{k+1}$, we deduce

$$(A.6) \quad du[0] \subset \mathcal{E}_{k+1} + \{du_j[0] | j \in J_k\}.$$

By the definition of I_{k+1} ,

$$(A.7) \quad \mathcal{E}_{k+1} = \mathcal{E}_k \oplus \{\{dy_{i_k}[k+1] | i_k \in I_{k+1}\}^*\} \oplus \{\{dy_\ell[k+1] | \ell \in I_{k+1} \setminus I_k\}^*\}.$$

For each $\ell \in I_{k+1} \setminus I_k$, $dy_\ell[k] \in \mathcal{E}_k$, and thus,

$$(A.8) \quad \begin{aligned} dy_\ell[k+1] &\in \{\{dx[1], dy_{i_0}[1], \dots, dy_{i_k}[k+1] | i_r \in I_r, 0 \leq r \leq k\}\} \\ &\subset (\mathcal{E}_k \oplus \{\{dy_{i_k}[k+1] | i_k \in I_k\}^*\}) + [du[0]] \\ &\subset (\mathcal{E}_k \oplus \{\{dy_{i_k}[k+1] | i_k \in I_k\}^*\}) + [du[0] | j \in J_k], \end{aligned}$$

where the last inclusion is from (A.3) for $t = k$. Thus, $(|I_{k+1}| - |I_k|)$ elements of $\{du_j[0] | j \in J_k\}$ are not independent of \mathcal{E}_{k+1} . One can therefore choose $J_{k+1} \subset J_k$ such that

$$(A.9) \quad du[0] \subset \mathcal{E}_{k+1} \oplus \{\{du_j[0] | j \in J_{k+1}\}^*\}$$

and

$$(A.10) \quad |J_{k+1}| = |J_k| - (|I_{k+1}| - |I_k|).$$

To finish up, from (A.5) and (A.9) it follows that

$$(A.11) \quad [dx, du[0], \dots, du[k+1]] \subset [\{dx, dy_{i_0}[0], \dots, dy_{i_{k+1}}[k+1], du_{j_{k+1}}[0], \dots, du_{j_0}[k+1] | i_t \in I_t, j_{k+1-t} \in J_t, 0 \leq t \leq k+1\}].$$

Since the reverse inclusion is obviously true, one has equality. By counting the number of vectors on the right-hand side of (A.11), we obtain (A.2) for $t = k + 1$. \square

From (A.1) and (A.2), respectively, it follows that

$$(A.12) \quad \rho_k = n + \sum_{i=1}^k |I_i|$$

and

$$(A.13) \quad n + (k+1)m = n + \sum_{i=1}^k |I_i| + \sum_{i=1}^k |J_i|,$$

yielding

$$(A.14) \quad \rho_k - \rho_{k-1} = m - |J_k|.$$

This combined with (A.3) proves the following result.

COROLLARY A.2. *Suppose that (2.1) is generically submersive. Then*

$$\dim \mathcal{E}_k - \dim \mathcal{E}_{k-1} = m \text{ if and only if } \{du_1[0], \dots, du_m[0]\} \subset \mathcal{E}_k .$$

Proof of Lemma 2.4. We can view $y[k], 0 \leq k \leq N$ as being an analytic function on $X \times U^{N+1}$. Let \mathcal{O} be an open subset of $X \times U^{N+1}$.

By the definition of I_0, \dots, I_n , for any $0 \leq N \leq n$,

$$(A.15) \quad dy_j[N] \in \text{span}_{\mathcal{K}_N} \{dx, dy_{i_0}[0], \dots, dy_{i_N}[N] \mid i_k \in I_k, 0 \leq k \leq N\} .$$

Due to analyticity, (A.15) is equivalent to

$$(A.16) \quad dy_j[N] \mid_{\mathcal{O}} \in \text{span}_{\mathbb{R}} \{dx, dy_{i_0}[0], \dots, dy_{i_N}[N] \mid i_k \in I_k, 0 \leq k \leq N\} \mid_{\mathcal{O}} ,$$

where the left-hand side is viewed as a one-form on \mathcal{O} , the right-hand side is viewed as an analytic codistribution on \mathcal{O} , and the span is taken pointwise; without loss of generality, it can be assumed that the codistribution has constant dimension. After possibly shrinking \mathcal{O} , the Rank theorem implies that on \mathcal{O} , $y_j[N]$ can be expressed as an analytic function of $(x, y_{i_0}[0], \dots, y_{i_N}[N] \mid i_k \in I_k, 0 \leq k \leq N)$. Repeating the above reasoning, (2.21) is equivalent to

$$(A.17) \quad dy_j[N] \mid_{\mathcal{O}} \in \text{span}_{\mathcal{K}_N} \{dy_j[0], \dots, dy_j[N-1], dy_{i_0}[0], \dots, dy_{i_N}[N] \mid i_k \in I_k, 0 \leq k \leq N\} \mid_{\mathcal{O}} .$$

Hence, from (A.16), (2.21) holds if

$$(A.18) \quad \frac{\partial y_j[N]}{\partial x} dx \in \text{span}_{\mathcal{K}_N} \left\{ \frac{\partial y_j[0]}{\partial x} dx, \dots, \frac{\partial y_j[N-1]}{\partial x} dx \right\} .$$

Because the right-hand side of (A.18) can be at most n -dimensional, there must exist an $N, 1 \leq N \leq n$, such that this is the case. \square

Appendix B.

Proofs of Lemmas 2.6 and 2.7. The inclusion $M_{k+1} \subset M_k$, implies $d_{k+1} \leq d_k \forall k \geq 0$. Since $x[k] : \mathbb{R}^n \times (\mathbb{R}^m)^k \rightarrow \mathbb{R}^n$ is an analytic function, there exists an open and dense subset $V_k \subset \mathbb{R}^n \times (\mathbb{R}^m)^k$ on which $x[k]$ has constant \mathbb{R} -rank equal to d_k ; that is, for each point $p \in V_k$,

$$(B.1) \quad \text{rank}_{\mathbb{R}} \left[\frac{\partial x[k]}{\partial(x, u[0], \dots, u[k-1])} \right] (p) = d_k .$$

Let $N_k := x[k](V_k)$. Then, N_k is an immersed submanifold of \mathbb{R}^n , but since the resulting topology may be different than the subset topology of \mathbb{R}^n , this property is of little interest. More importantly, the implicit function theorem implies that, for each point $q \in N_k$, there exists an open subset $\mathcal{O}_k \subset V_k$ such that $x[k](\mathcal{O}_k)$ is a d_k -dimensional, embedded, analytic submanifold of \mathbb{R}^n and q is in the interior of $x[k](\mathcal{O}_k)$. Since V_k is dense in $\mathbb{R}^n \times (\mathbb{R}^m)^k$, N_k is dense in M_k in the subset topology.

Let $\bar{f}_k : x[k](\mathcal{O}_k) \times \mathbb{R}^m \rightarrow \mathbb{R}^n$ denote f restricted to $x[k](\mathcal{O}_k) \times \mathbb{R}^m$ (the subscript k is to note that \bar{f}_k depends on \mathcal{O}_k). Whenever $\mathcal{O}_k \subset V_k$ is such that $x[k](\mathcal{O}_k)$ is a d_k -dimensional embedded submanifold of \mathbb{R}^n , then \bar{f}_k is an analytic function. Consequently, it will have constant \mathbb{R} -rank on an open and dense subset of its domain

of definition, which is called its *generic rank* and is denoted as $\text{gen rank } \bar{f}_k$. From (2.23) and (2.24) it follows that

$$(B.2) \quad d_{k+1} = \text{gen rank } \bar{f}_k : x[k](\mathcal{O}_k) \times \mathbb{R}^m \rightarrow \mathbb{R}^n$$

since, if $p \in \mathcal{O}_k$ and $q := x[k](p)$, then $T_q x[k](\mathcal{O}_k)$, the tangent space of $x[k](\mathcal{O}_k)$ at the point q , satisfies

$$(B.3) \quad T_q x[k](\mathcal{O}_k) = \text{Image} \left[\frac{\partial x[k]}{\partial (x, u[0], \dots, u[k-1])} \right] (p).$$

With the above preliminaries completed, the proof of Lemma 2.6 can be given. If $d_n = 0$, the result is obvious; suppose, therefore, that $d_n > 0$. Then, there exists $0 \leq k \leq n-1$ such that $d_k = d_{k-1}$. As before, choose \mathcal{O}_k to be an open subset of V_k such that $x[k](\mathcal{O}_k)$ is an embedded d_k -dimensional submanifold of \mathbb{R}^n . Since $M_k \subset M_{k-1}$, and N_k and N_{k-1} are dense in M_k and M_{k-1} , respectively, the condition $d_k = d_{k-1}$ implies that $x[k](\mathcal{O}_k)$ is an embedded $(d_{k-1} = d_k)$ -dimensional submanifold of \mathbb{R}^n and $x[k](\mathcal{O}_k) \cap x[k-1](\mathcal{O}_{k-1})$ has nonempty interior. Therefore,

$$(B.4) \quad \begin{aligned} d_{k+1} &= \text{gen rank } \bar{f}_k : x[k](\mathcal{O}_k) \times \mathbb{R}^m \rightarrow \mathbb{R}^n \\ &= \text{gen rank } \bar{f}_{k-1} : x[k-1](\mathcal{O}_{k-1}) \times \mathbb{R}^m \rightarrow \mathbb{R}^n \\ &= d_k, \end{aligned}$$

where the fact that \bar{f}_k and \bar{f}_{k-1} are the restrictions of a common map f and have a common nonempty open set in their domain of definition entails the second equality. This completes the proof of Lemma 2.6.

Turning to Lemma 2.7, let $d := d_n$, which is then equal to d_k for all $k > n$ by Lemma 2.6. If $d = 0$, then Lemma 2.7 is immediate, so in the following it is supposed that $d > 0$. Let $\alpha = \alpha(x, u[0], \dots, u[k])$ be an element of \mathcal{R}_k . For any $r > 0$, $\delta^r(\alpha) = \alpha(x[r], u[r], \dots, u[k+r])$. Let $\tilde{\mathcal{O}}_r \subset V_r$ be an open set such that $x[r](\tilde{\mathcal{O}}_r)$ is a d_r -dimensional submanifold of \mathbb{R}^n . Then $\delta^r(\alpha) \neq 0$ if, and only if, α restricted to $(x[r](\tilde{\mathcal{O}}_r) \times (\mathbb{R}^m)^{k+1}) \neq 0$. When $r > n$, there exists $\tilde{\mathcal{O}}_n \subset V_n$ such that $x[n](\tilde{\mathcal{O}}_n) \times (\mathbb{R}^m)^{k+1} \cap x[r](\tilde{\mathcal{O}}_r) \times (\mathbb{R}^m)^{k+1}$ has nonempty interior. Thus, α restricted to $(x[r](\tilde{\mathcal{O}}_r) \times (\mathbb{R}^m)^{k+1}) \neq 0$ if, and only if, α restricted to $(x[n](\tilde{\mathcal{O}}_n) \times (\mathbb{R}^m)^{k+1}) \neq 0$. \square

Acknowledgment. The author has benefited greatly from conversations with M. Fliess and C.H. Moog on the subject of this paper, as well as from the insightful comments provided by the reviewers. This work was completed while the author was a visiting researcher at the Laboratoire des Signaux et Systèmes, Gif-sur-Yvette, France (Poste Rouge, Directeur de Recherche Associé). Professors P. Bertrand, M. Fliess, and F. Lamnabhi-Lagarrigue are sincerely thanked for contributing in diverse ways to a very enjoyable and worthwhile *séjour*.

REFERENCES

- [1] LI CAO AND YU-FAN ZHENG, *On minimal compensators for decoupling control*, preprint, 1991.
- [2] M. D. DI BENEDETTO AND J. W. GRIZZLE, *An analysis of regularity conditions in nonlinear synthesis problems*, Lecture Notes in Control and Inform. Sci., Vol. 144, Springer-Verlag, Berlin, 1990, pp. 843-850.
- [3] ———, *Intrinsic notions of regularity for local inversion, output nulling and dynamic extension of non-square systems*, Control Theory Adv. Tech., 6 (1990), pp. 357-381.

- [4] M. D. DI BENEDETTO, J. W. GRIZZLE, AND C. H. MOOG, *Rank invariants of nonlinear systems*, SIAM J. Control Optim., 72 (1989), pp. 658–672.
- [5] S. DIOP AND M. FLIESS, *Nonlinear observability, identifiability and persistent trajectories*, in Proc. 30th IEEE Conference on Decision and Control, Brighton, England, December 1991, pp. 714–718.
- [6] S. EL ASMI AND M. FLIESS, *Formules d'inversion*, in Analysis of Controlled Dynamical Systems, B. Bonnard, B. Bride, J.P. Gauthier, and I. Kupka, eds., Conf. Proc., Lyon, France, July 1990, Birkhäuser, Boston, MA.
- [7] ———, *Inversion formula for discrete-time systems*, IFAC Symposium on Nonlinear Control Systems Design, NOLCOS-92, Bordeaux, France, June 1992.
- [8] M. FLIESS, *A new approach to the noninteracting control problem in nonlinear systems theory*, in Proc. 23rd Allerton Conference, University of Illinois, Monticello, IL, 1985, pp. 123–129.
- [9] ———, *Generalized controller canonical forms for linear and nonlinear dynamics*, IEEE Trans. Automat. Control, AC-35, 9 (1990), pp. 994–1001.
- [10] ———, *Automatique en temps discret et algèbre aux différences*, Forum Mathematicum, 2 (1990), pp. 213–232.
- [11] ———, *Reversible linear and nonlinear discrete time dynamics* IEEE Trans. Automat. Control, AC-37, (1992), pp. 1144–1153.
- [12] M. FLIESS AND D. NORMAND-CYROT, *A group theoretic approach to discrete-time nonlinear controllability*, Proc. 20th IEEE Conf. Decision and Control, San Diego, 1981, pp. 551–557.
- [13] A. GLUMINEAU AND C. H. MOOG, *Essential orders and the nonlinear decoupling problem*, Internat. J. Control, 50 (1989), pp. 1825–1834.
- [14] J. W. GRIZZLE AND H. NIJMEIJER, *Zeros at infinity for nonlinear discrete-time systems*, Math. Syst. Theory, 19 (1986), pp. 79–93.
- [15] R. HERMAN AND H. K. KRENER, *Nonlinear controllability and observability*, IEEE Trans. Automat. Control, 22 (1977), pp. 728–740.
- [16] A. J. C. HUIJBERTS, H. NIJMEIJER, AND L. L. M. VAN DER WEGEN, *Dynamic disturbance decoupling for nonlinear systems*, SIAM J. Control Optim., 30 (1992), pp. 336–349.
- [17] B. JAKUBCZYK AND E. D. SONTAG, *Controllability of nonlinear discrete-time system: a Lie algebraic approach*, SIAM J. Control Optim., 28 (1990), pp. 1–33.
- [18] Ü. KOTTA, *Right-inverse of a discrete-time non-linear system*, Internat. J. Control, 51 (1990), pp. 1–9.
- [19] ———, *Local (finite-time) model matching of nonlinear discrete-time systems*, preprint, 1990.
- [20] D. LUENBERGER, *Optimization by Vector Space Methods*, John Wiley, New York, 1969.
- [21] J. L. MASSEY AND M. K. SAIN, *Invertibility of linear time-invariant dynamic systems*, IEEE Trans. Automat. Control, AC-14 (1969), pp. 141–149.
- [22] C. H. MOOG, *Nonlinear decoupling and structure at infinity*, Math. Control Signals Systems, 1 (1988), pp. 257–268.
- [23] H. NIJMEIJER, *Right-invertibility for a class of nonlinear control systems: a geometric approach*, Systems Control Lett., 7 (1986), pp. 125–132.
- [24] ———, *On dynamic decoupling and dynamic path controllability in economic systems*, J. Econom. Dynamics Control, 13 (1989), pp. 21–39.
- [25] D. NORMAND-CYROT, *Théorie and pratique des systèmes non linéaires en temps discret*, Thèse d'Etat, Université Paris-Sud, Orsay, 1983.
- [26] L. M. SILVERMAN, *Inversion of multivariable linear systems*, IEEE Trans. Automat. Control, AC-14 (1969), pp. 270–276.
- [27] S. N. SINGH, *A modified algorithm for invertibility in nonlinear systems*, IEEE Trans. Automat. Control, AC-26 (1981), pp. 595–598.
- [28] E. D. SONTAG, *Polynomial Response Maps*, Lecture Notes in Control and Inform. Sci., 13, Springer, Berlin, 1979.
- [29] E. D. SONTAG AND Y. ROCHALEAU, *On discrete-time polynomial systems*, Nonlinear Anal.: Theory Meth. and Appl., 1 (1976), pp. 55–64.
- [30] B. L. VAN DER WAERDEN, *Modern Algebra*, Frederick Ungar Pub., New York, Vols. I and II, 1953.
- [31] Y. WANG AND E. D. SONTAG, *Realizations and I/O relations: The analytic case*, Proc. IEEE Conference on Decision and Control, Tampa, December 1989, pp. 1275–1980.
- [32] X.-H. XIA, *The essential structure of nonlinear systems*, preprint, 1989.